

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/005500 A1

(51) International Patent Classification: C12N 9/10

(21) International Application Number:
PCT/KR2003/001301

(22) International Filing Date: 2 July 2003 (02.07.2003)

(25) Filing Language: Korean

(26) Publication Language: English

(30) Priority Data:
10-2002-0038011 2 July 2002 (02.07.2002) KR

(71) Applicants (for all designated States except US):
GENOMINE INC. [KR/KR]; Environmental Eng. Bldg.
226, Pohang University of Science and Technology, Po-
hang, Kyungbuk 790-784 (KR). KOREA RESEARCH
INSTITUTE OF CHEMICAL TECHNOLOGY
[KR/KR]; P.O. Box 107, Yusong, Taejeon 305-600 (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LEE, Dong-Hee
[KR/KR]; 175-20 Yconji-dong, Busanjin-gu, Busan

614-070 (KR). HWANG, In-Taek [KR/KR]; Han-
vit Apt. 115-303, Eeun-dong, Yusong-gu, Taejeon
305-755 (KR). CHOI, Jung-Sup [KR/KR]; Hanwool
Apt. 102-1106, Sinseong-dong, Yusong-gu, Taejeon
305-345 (KR). MIN, Yong-Ki [KR/KR]; Hwangsil Apt.
117-106, Wolpyeong-dong, Seo-gu, Taejeon 302-280
(KR). KIM, Tae-Joon [KR/KR]; Expo Apt. 305-302g,
Busanjin-gu, Jeonmin-dong, Yusong-gu, Taejeon 305-761
(KR). KO, Jae-Heung [KR/KR]; Kwangsung Villa 401,
214, Mapo-dong, Mapo-gu, Seoul 121-050 (KR). KIM,
Tae-Hoon [KR/KR]; 264-8 Bugok-dong, Geumjeong-gu,
Busan 609-320 (KR). PARK, Yu-Shin [KR/KR]; Seungri
Apt. 2-403, 253-6, Hyoja-dong, Nam-gu, Pohang-si,
Gyeongsangbuk-do 790-330 (KR). CHO, Kwang-Yun
[KR/KR]; 383-21 Doryong-dong, Yusong-gu, Taejeon
305-340 (KR). LEE, Seon-Woo [KR/KR]; Hanvit Apt.
103-805, Eeun-dong, Yusong-gu, Taejeon 305-755 (KR).

(74) Agent: KIM, Seog-Hyun; 9th Floor, Daekyung Bldg.
120, 2-ka, Taepyung-ro, Chung-ku, Seoul 100-724 (KR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

[Continued on next page]

(54) Title: NOVEL POLYPEPTIDE HAVING FUNCTION OF 7-KETO-8-AMINOPELARGONIC ACID SYNTHASE OF
PLANT AND METHOD FOR INDUCING GROWTH INHIBITION AND LETHALITY BY SUPPRESSING EXPRESSION
OF THE POLYPEPTIDE



(57) Abstract: The invention relates to a novel polypeptide participating in biotin biosynthesis of plant, a polynucleotide encoding the said polypeptide, a method for inducing plant growth inhibition by suppressing the expression or function of the said polypeptide, resulting in inhibition of biotin biosynthesis, and a method for identifying herbicidal compounds that inhibit the expression or function of the said polypeptide. Biotin biosynthesis is an essential process for the growth of plants, however, the process is not present in both human and animals. Therefore, the compounds suppressing the expression or function of the novel polypeptide that relates to biotin biosynthesis can be effectively used as a new herbicide doing human and animals no harm.

WO 2004/005500 A1

BEST AVAILABLE COPY

WO 2004/005500 A1



CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,
US, UZ, VC, VN, YU, ZA, ZM, ZW.

ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

- (84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/005500

PCT/KR2003/001301

NOVEL POLYPEPTIDE HAVING FUNCTION OF 7-KETO-8
- AMINOPELARGONIC ACID SYNTHASE OF PLANT AND
METHOD FOR INDUCING GROWTH INHIBITION AND
LETHALITY BY SUPPRESSING EXPRESSION OF THE
POLYPEPTIDE

5

FIELD OF THE INVENTION

The present invention relates to a novel polypeptide isolated from plant and
a polynucleotide encoding the same, more particularly a novel polypeptide involved
in biotin biosynthesis of plant, a polynucleotide encoding the said polypeptide, a
method for inducing plant growth inhibition by suppressing expression or function of
the said polypeptide, resulting in inhibition of biotin synthase activity, and a method
for identifying herbicidal compounds that inhibit the expression or the function of the
said polypeptide.

15

BACKGROUND OF THE INVENTION

As the most recently found member of vitamin B group, biotin is essential
for the growth of both animals and plants. Biotin plays an important role as a
coenzyme in transporting CO₂ to a specific substrate during carboxylation,
decarboxylation and transcarboxylation related to fatty acid or carbohydrate
metabolism. Bacteria, plants and some of algae can synthesize biotin *in vivo*, while
most algae and animals cannot, suggesting that they should get biotin from outside.

25

Since the biotin biosynthesis pathway was first disclosed in *Escherichia coli*

WO 2004/005500

PCT/KR2003/001301

and *Bacillus subtilis* (Eisenberg MA, *Adv. Enzymol.*, 38:317-372, 1973; Pai CH, *J. Bacteriol.*, 121:1-8, 1975), studies have been actively made on the biotin biosynthesis pathway of bacteria based on biochemistry and genetics (Ploux and Marquet, *Biochem. J.*, 283:327-331, 1992; Alexeev *et al.*, *J. Mol. Biol.*, 235:774-776, 1994; Huang *et al.*, *Biochemistry*, 34:10985-10995, 1995).

As a result, which gene plays what role in each stage of biotin biosynthesis pathway in microorganisms such as *Escherichia coli* and *Bacillus subtilis* has been clearly answered. Particularly, it has been confirmed that *Escherichia coli* has a *bio* cluster consisting of 5 genes of *bioABFCD* and *Bacillus subtilis* has a *bio* cluster consisting of 6 genes of *bioWAFDBI* (Bachman BJ, *Microbiol. Rev.*, 54:130-197, 1990; Bower *et al.*, *J. Bacteriol.*, 178:4122-4130, 1996). *Bacillus sphearicus* was confirmed to have two independent clusters, *bioXWF* and *bioDAYB*. In addition, it was confirmed that the expression of biotin in *Escherichia coli* is suppressed by the attachment of BirA protein to the region between *bioA* and *bioB* genes (Barker and Campbell, *J. Mol. Biol.*, 146:469-492, 1981).

The first step of biotin biosynthesis in *Escherichia coli* is decarboxylative condensation catalyzed by KAPA synthase (7-keto-8-aminopelargonic acid synthase), in which L-alanine and pimeloyl CoA were converted into KAPA (7-keto-8-aminopelargonic acid, also known as 8-amino-7-oxononanoate; AON). KAPA synthase, a *bioF* gene product of *Escherichia coli*, is a pyridoxal 5'-phosphate-dependent enzyme and has about 42 kDa molecular weight.

As seen hereinbefore, biotin biosynthesis pathway of microorganism has been well known, while that in plants has not been explained yet, especially there are much to be known about KAPA synthase playing a role in the early stage of biotin biosynthesis. However, some interesting evidence has been reported that biotin biosynthesis is occurring in plants following the same pathways as in *Escherichia*

WO 2004/005500

PCT/KR2003/001301

coli (Baldet *et al.*, *Eur. J. Biochem.*, 217:479-485, 1993). Moreover, biotin biosynthesis in plants and the use thereof have also been disclosed by the research on biotinylated protein (Nikolau *et al.*, *Anal. Biochem.*, 149:448-453, 1985; Tissot *et al.*, *Biochem. J.*, 314:391-395, 1996), and the isolation and the research on the characteristics of auxotrophic mutants. Particularly, the detailed analysis on mutants like auxotrophic mutants offered great information about biotin biosynthesis and its regulation.

Firstly, converting reaction from KAPA to 7,8-diaminopelargonic acid (DAPA) is absent in *bio1* auxotrophic mutant of *Arabidopsis thaliana* (Meinke DW, *Theor. Appl. Genet.*, 72:543-552, 1985). And the last step of biotin biosynthesis pathway, which is converting reaction from dethiobiotin to biotin, is absent in *bio2* mutant (Patton *et al.*, *Plant Physiol.*, 116:935-946, 1998).

The above studies support the possibility to develop an environmental friendly herbicide based on the study on biotin biosynthesis in plants. Biotin biosynthesis is indispensable procedure for plant growth, but the process is not present in both human and animals. Thus, it could be possible to inhibit the growth of harmful plants by suppressing biotin biosynthesis, without doing harm to human and animals. Rendina *et al.* tried to develop a new herbicide by targeting bacterial dethiobiotin synthetase (DTBS) (Rendina *et al.*, *Pesticide Sci.*, 55:236-247, 1999). But, as of today, biotin biosynthesis in plants has not been fully explained and the studies on a method to inhibit the growth of harmful plants by targeting genes related to biotin biosynthesis, which seems to be safe and environmental friendly, have just begun. And the functions of metabolism in plants have not been clearly disclosed when the conventional herbicides are used.

25

WO 2004/005500

PCT/KR2003/001301

DISCLOSURE OF THE INVENTION

Thus, the present inventors have studied on the biotin biosynthesis pathway taking *Arabidopsis thaliana* as a major subject. As a result, the inventors isolated a polypeptide participated in the first step of biotin biosynthesis pathway and a polynucleotide encoding the same, and analyzed the functions thereof. And also, the inventors confirmed that the suppression of the expression or function of the said polypeptide could be a fatal blow for plant growth. The present inventors have completed this invention by confirming that the said polynucleotide and the polypeptide expressed therefrom could be excellent candidates for developing novel herbicides.

It is an object of the present invention to provide a method for the development of a novel environmental friendly herbicide inducing plant growth inhibition by suppressing biotin biosynthesis in plants.

In order to achieve the above object, the present invention provides an isolated polypeptide having an amino acid sequence selected from the group consisting of;

- (a) an amino acid sequence of SEQ. ID. NO. 2, and
- (b) an amino acid sequence having at least 70% identity to SEQ. ID. NO. 2.

The present invention also provides an isolated polynucleotide encoding the said polypeptide or an antisense polynucleotide complementary to the same.

The present invention provides an expression vector comprising the isolated polynucleotide encoding the said polypeptide or the antisense polynucleotide complementary to the same.

WO 2004/003500

PCT/KR2003/001301

The present invention provides a cell comprising the said expression vector.

The present invention provides a transgenic plant or seed comprising the said expression vector.

The present invention further provides a method for inducing plant growth inhibition by suppressing the expression or the function of the said polypeptide, resulting in the inhibition of biotin biosynthesis..

The present invention yet further provides a method for identifying herbicidal compounds that inhibit the expression or function of the said polypeptide.

10 In the present invention, the gene encoding KAPA synthase of *Arabidopsis thaliana* is stated in italics such as "*AtKAPAS* polynucleotide", "*AtKAPAS* gene" or "*AtKAPAS*" and the protein encoded thereby is stated as "AtKAPAS polypeptide", "AtKAPAS protein" or "AtKAPAS".

15 Further features of the present invention will appear hereinafter.

The present invention provides a novel polypeptide involved in biotin biosynthesis. The polypeptide has the same function as *E. coli* KAPA synthase that plays a role in converting L-alanine and pimeloyl CoA into KAPA (7-keto-8-aminopelargonic acid) in the first step of biotin biosynthesis pathway in *E. coli*.

20 The range of polypeptide according to the present invention may include a polypeptide having an amino acid sequence represented by SEQ ID No. 2, and functional equivalents thereof. The "functional equivalents" refer to polypeptides having more than 70%, preferably more than 80%, more preferably more than 90% identity (sequence homology) to the amino acid sequence of SEQ ID No. 2 as a result of addition, substitution or deletion of one or more amino acids, and exhibiting substantially the same physiological activity of the polypeptide of SEQ ID No. 2.

WO 2004/005500

PCT/KR2003/001301

Also, "substantially the same physiological activity" means an activity converting L-alanine and pimeloyl CoA into KAPA (7-keto-8-aminopelargonic acid) in the first step of biotin biosynthesis pathway. The protein according to the present invention can be obtained from nature (for example, plant cells) or by expression of a recombinant nucleic acid that encodes the protein or by a chemical synthesis. Preferably, the protein can be isolated from *Arabidopsis thaliana*.

The present invention also provides a polynucleotide encoding the said polypeptide. More particularly, the present invention provides an *AtKAPAS* (*Arabidopsis thaliana* KAPA synthase), a polynucleotide having a nucleotide sequence represented by SEQ. ID. NO. 1. The *AtKAPAS* has about 51.3 kDa molecular weight and comprises 1410 bp size open reading frame encoding 469 amino acids.

The present invention provides an antisense polynucleotide complementary to the said polynucleotide. More particularly, the present invention provides an antisense polynucleotide complementary to the polynucleotide having a nucleotide sequence of SEQ. ID. NO. 1. Antisense polynucleotides are generally known to combine with target nucleotides in nucleic acid (RNA or DNA), by which they suppress the synthesis or function of the nucleic acid. Particularly, an antisense polynucleotide corresponding to a specific gene can be hybridized with both RNA and DNA, leading to the suppression of the expression of the specific gene in the level of transcription or translation.

Meanwhile, the amino acid sequence encoded by the *AtKAPAS* shows 28%, 34% and 38% homology and 43%, 52% and 54% similarity, respectively, to the sequences of *bioF* genes (GeneBank accession number NP286539, NP390900 and

WO 2004/005500

PCT/KR2003/001301

JQ0512, each) encoding KAPA synthase, a protein involved in biotin biosynthesis of *Escherichia coli*, *Bacillus subtilis* and *Bacillus sphaericus*. The amino acid sequence also comprises aminotransferase class I and class II domain at its C-terminal, suggesting that the polynucleotide of the invention has the function of
5 aminotransferase in addition to that of KAPA synthase. Besides, the amino acid sequence comprises a domain presumed to be plasma membrane spanning region.

The polynucleotide of the present invention that encodes the polypeptide involved in biotin synthesis or the antisense polynucleotide that is complementary
10 thereto can be inserted to a suitable expression vector to transform plant cells. The "expression vector" refers to a known plasmid, virus or other medium into which the polynucleotide sequence encoding the polypeptide involved in biotin synthesis or the antisense polynucleotide complementary thereto can be inserted or introduced. The gene sequence of the present invention can be operably linked to the expression
15 control sequence. The operably linked gene sequence and expression control sequence can be included within a single expression vector containing a selective marker and a replication origin. The "operably linked gene sequence and expression control sequence" may be a gene sequence and an expression control sequence which are linked in such a manner to enable the expression of the gene
20 when a suitable molecule is bound to the expression control sequence. The "expression control sequence" means a DNA sequence controlling the expression of a nucleic acid sequence which is operably linked in a particular host cell. Such a control sequence includes a promoter for performing transcription, an operator sequence for controlling transcription, a sequence encoding a suitable mRNA
25 ribosome binding site, and a sequence controlling termination of transcription or translation. A suitable vector into which the gene of the present invention can be

WO 2004/005500

PCT/KR2003/001301

introduced is a Ti plasmid, a root inducing (Ri) plasmid or a plant virus vector. The suitable vector may be, but not limited to, a binary vector of pCAL-n or pCAMBIA series. Anyone skilled in the pertinent art can select a suitable vector for introducing the nucleic acid of the gene of the present invention. Any vector
5 capable of introducing the polynucleotide sequence encoding the polypeptide involved in biotin synthesis or the antisense polynucleotide complementary thereto into plant cells can be used in the present invention.

The expression vector according to the present invention can be introduced into cells by known methods. The cells may be eukaryotic cells such as yeast or
10 plant cell, or prokaryotic cells, preferably *E. coli* or *Agrobacterium* sp. The methods for introducing the expression vector of the present invention to host cells include, but not limited to transformation using *Agrobacterium* species microorganisms, particle gun bombardment, silicon carbide whiskers, sonication, electroporation and PEG (polyethyleneglycol) precipitation. The method may be
15 preferably the transformation using *Agrobacterium* species microorganisms or electroporation. Accordingly, the present invention provides a cell transformed with the expression vector of the present invention. The host cell includes eukaryotic cells such as yeast or plant cell, or prokaryotic cells such as *E. coli*.
Beside, the present invention provides plants to which the expression vector was
20 introduced and seeds obtained from the plants.

In the preferred embodiments of the present invention, the inventors constructed pCKAPA recombinant vector by introducing *AtKAPAS* cDNA into pCAL-n vector (Stratagene, USA) and prepared *E. coli* transformed with the
25 recombinant vector, which was deposited at Korean Collection for Type Cultures of Korea Research Institute of Biotechnology and Bioscience on March 26, 2002

WO 2004/005500

PCT/KR2003/001301

(Accession No: KCTC 10210BP). Since the above pCAL-n vector contains calmodulin-binding peptide tag sequence, the protein expressed by the vector can be easily separated by calmodulin resin.

5 The present invention further provides a method for inducing plant growth inhibition by suppressing the expression or function of the polypeptide involved in biotin synthesis, resulting in inhibition of biotin biosynthesis.

 In order to suppress the expression of the said polypeptide, conventional methods generally known to those skilled the pertinent art are all available, that is,
10 antisense polynucleotide introduction, gene deletion, gene insertion, T-DNA introduction, homologous recombination or transposon tagging, etc., and particularly, the method to introduce antisense polynucleotide into plants was used in this invention. The present inventors prepared an antisense construct containing antisense polynucleotide sequence of *AtKAPAS*, a polynucleotide having a nucleotide
15 sequence represented by SEQ. ID. NO. 1. Precisely, the inventors constructed pSEN-K recombinant vector by introducing a polynucleotide having a nucleotide sequence represented by SEQ. ID. NO. 1 into pSEN vector in the antisense orientation and transformed *E. coli* with the prepared recombinant vector. The *E. coli* comprising the pSEN-K recombinant vector was deposited at Korean Collection
20 for Type Cultures of Korea Research Institute of Biotechnology and Bioscience on March 26, 2002 (Accession No: KCTC 10211BP).

 The function of the polypeptide according to the present invention is preferably suppressed by treatment of chemical substances. As a chemical
25 substance used for suppressing the function of the polypeptide, pimeloyl CoA derivatives working as a substrate to KAPA synthase in the biotin biosynthesis

WO 2004/005500

PCT/KR2003/001301

pathway can be selected. As stated in an embodiment of the present invention, the polypeptide expressed from the *AtKAPAS* polynucleotide has strong substrate-specificity especially to pimeloyl CoA. Therefore, pimeloyl CoA derivatives working as competitive inhibitors with pimeloyl CoA can effectively inhibit biotin biosynthesis in plants, and thus, they can be used for inducing plant growth inhibition.

In another embodiment of the present invention, the inventors inhibited protein synthesis from *AtKAPAS* by transforming *Arabidopsis thaliana* with the pSEN-K recombinant vector. As a result, it was confirmed that serious retardation of plant growth, leaf etiolation and plant death were induced by the antisense construct of *AtKAPAS*. Therefore, based on the fact that the antisense construct effectively inhibits plant growth, the present inventors confirmed that the polynucleotide of the present invention could be effectively used for the development of a novel herbicide.

Accordingly, the present invention provides a method for identifying herbicidal compounds that inhibit the expression or function of the polypeptide involved in biotin synthesis. Particularly, the present invention provides a method of identifying herbicidal compounds that inhibit the expression or function of the polypeptide of claim 1, comprising the steps of;

(a) combining a polypeptide of claim 1 with the compounds to be tested for the ability to inhibit the expression or the function of the polypeptide under conditions conducive to inhibition;

(b) selecting the identified compounds to inhibit the expression or function of the polypeptide in step (a);

(c) applying the said compounds selected in step (b) to a plant to test for herbicidal activity; and

WO 2004/005500

PCT/KR2003/001301

(d) selecting the identified compounds to have herbicidal activity in step (c).

The term "herbicidal compounds" refer to substances used to kill or suppress the growth of plants, plant cells, plant seeds, or plant tissues. The effect of herbicidal compounds on expression or function of the polypeptide according to the present invention can be identified by northern blot analysis and western blot analysis that are known to those skilled in the pertinent art. The herbicidal compounds may be, for example, peptides, polypeptides, peptide imitators, chemical compounds, biologicals and the like.

10

BRIEF DESCRIPTION OF THE DRAWINGS

The application of the preferred embodiments of the present invention is best understood with reference to the accompanying drawings, wherein:

15

FIG. 1 is a diagram showing the putative amino acid sequence encoded by the nucleotide sequence of *Arabidopsis thaliana* AtKAPAS gene, compared with KAPA synthase amino acid sequences of *Escherichia coli*, *Bacillus subtilis* and *Bacillus sphaericus*. Marked similar amino acid sequence regions with black blocks;

20

FIG. 2 is a graph showing the protein content in each fraction purified by calmodulin affinity chromatography using 2 mM EDTA from crude extracts purified from *E. coli* transformed with recombinant vector containing AtKAPAS gene and control *E. coli* transformed with a vector only;

25

● : *E. coli* transformed with recombinant vector containing AtKAPAS gene,

○ : Control *E. coli* transformed with a vector only

WO 2004/005500

PCT/KR2003/001301

FIG. 3 is a photograph showing the result of SDS-PAGE electrophoresis on fractions #12-14 among every chromatography fraction obtained from effluents of both *E. coli* transformed with recombinant vector containing *AtKAPAS* gene and control *E. coli*;

5 Lane 1: Fraction #12 of effluents obtained from *E. coli* transformed with recombinant vector containing *AtKAPAS* gene,

Lane 2: Fraction #12 of effluents obtained from control *E. coli*,

Lane 3: Fraction #13 of effluents obtained from *E. coli* transformed with recombinant vector containing *AtKAPAS* gene,

10 Lane 4: Fraction #13 of effluents obtained from control *E. coli*,

Lane 5: Fraction #14 of effluents obtained from *E. coli* transformed with recombinant vector containing *AtKAPAS* gene,

Lane 6: Fraction #14 of effluents obtained from control *E. coli*

FIG. 4 is a graph showing the Lineweaver-Burk plot reflecting the changes
15 of reaction speed according to the changes of substrate concentration, which was available for the measurement of the substrate-specific enzyme activity of the purified protein to pimeloyl CoA;

FIG. 5 is a diagram showing the structure of a vector into which *AtKAPAS* gene was introduced in the antisense orientation;

20 FIG. 6A is a photograph showing saplings growing up from seeds of transgenic *Arabidopsis thaliana* to which *AtKAPAS* gene was introduced in the antisense orientation (the saplings were grown and differentiated in a pot with biotin);

FIG. 6B is a photograph showing saplings growing up from seeds of
25 transgenic *Arabidopsis thaliana* to which *AtKAPAS* gene was introduced in the antisense orientation (the saplings were grown and differentiated in a pot without

WO 2004/005500

PCT/KR2003/001301

biotin).

BEST MODE FOR CARRYING OUT THE INVENTION

5 Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

10

Example 1: Isolation of *E. coli bioF* homologous gene from *Arabidopsis thaliana*

The present inventors performed screening to isolate homologous genes corresponding to *E. coli bioF*, a gene encoding KAPA synthase of biotin biosynthesis pathway, from *Arabidopsis thaliana*.

15

<1-1> *Arabidopsis thaliana* cultivation and culture

The present inventors cultured *Arabidopsis thaliana* either in pots filled with soil or in petri dishes containing MS (Murashige and Skoog salts, Sigma, USA) medium supplemented with 2% sucrose (pH 5.7) and 0.8% agar. Put the pots in
20 growth chambers where the temperature was kept at 22 °C and the light and dark cycle was set to 16/8 hours.

<1-2> RNA Extraction and cDNA library preparation

In order to construct *Arabidopsis thaliana* cDNA library, the total RNA was
25 extracted from *Arabidopsis thaliana* leaves in various differentiation stages using

WO 2004/005500

PCT/KR2003/001301

TRI reagent (Sigma, USA). Poly(A)+ RNA was isolated from the extracted total RNA according to the protocol of mRNA isolation kit (Pharmacia, USA). Double stranded cDNA was prepared with poly(A)+ RNA and cDNA synthesis kit (Time Saver cDNA synthesis kit, Pharmacia, USA) using *NotI*-(dT)18 primer.

5

<1-3> Isolation of *bioF* homologous gene

Based on the nucleotide sequence of *E. coli bioF* gene, the present inventors synthesized a forward primer (SEQ. ID. NO. 3) including the sequence of restriction enzyme *Bam*HI and a reverse primer (SEQ. ID. NO. 4) including the sequence of restriction enzyme of *Hind*III. By performing PCR (polymerase chain reaction) with the two primers, the full-length cDNA was amplified and isolated from *Arabidopsis thaliana* cDNA library prepared in the above Example <1-2>.

From the sequence analysis of the isolated cDNA, it was confirmed that the cDNA has about 51.3 kDa molecular weight and contains 1410 bp size open reading frame encoding 469 amino acids, and the cDNA was named *AtKAPAS* (*Arabidopsis thaliana* KAPA synthase).

The putative amino acid sequence encoded by the *AtKAPAS* showed 28%, 34% and 38% homology and 43%, 52% and 54% similarity to the sequences of *bioF* genes (GeneBank accession number NP286539, NP390900 and JQ0512, each) of *Escherichia coli*, *Bacillus subtilis* and *Bacillus sphaericus* (see FIG. 1), respectively.

Besides, it was confirmed that the above protein contains aminotransferase class I and class II domain at C-terminal region, and a domain predicted to be the plasma membrane spanning region.

WO 2004/005500

PCT/KR2003/001301

Example 2: Purification of protein expressed by *AtKAPAS* gene in *E. coli*

<2-1> Induction of protein expression

The amplified DNA fragment comprising the full-length *AtKAPAS* cDNA isolated in the Example <1-3> was cut with restriction enzymes of *Bam*HI and *Hind* III, and then constructed pCKAPA recombinant vector by cloning the fragment in the region of *Bam*HI and *Hind*III site of pCAL-n vector (Stratagene, USA).

E. coli BL21-Gold(DE)(Stratagene, USA) was transformed with the pCKAPA recombinant vector (Accession No: KCTC 10210BP), and then cultured in LB (Luria-Bertani broth, USB, USA) medium containing 100 µg/ml of ampicillin at 37°C with agitation (150 rpm) until the value of OD₆₀₀ reached 0.7. In order to induce the expression of the target protein in *E. coli* cells, IPTG (isopropyl-D-thiogalactoside) was added to the suspension at a final concentration of 1mM, and further cultured for 2 more hours. The culture cells were washed with 50 mM potassium phosphate buffer (pH 7.0) containing 50 mM MgSO₄ and 0.4 M NaCl, after which centrifuged at 4,000× g for 15 minutes. The precipitates thereof were collected and stored at -20°C.

<2-2> Protein purification

The cell precipitates obtained in the above Example <2-1> were suspended in CaCl₂ binding buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 10 mM β-mercaptoethanol, 1.0 mM magnesium acetate, 1.0 mM imidazole, 2 mM CaCl₂). Lysozyme was added into the cell suspension at a final concentration of 200 µg/ml and the cell suspension was stirred for 15 minutes. Then, ultrasonication was performed for 30 seconds. The crushed samples were cooled down with ice for 5 minutes. Such procedure (cooling after ultrasonication) was repeated three times.

WO 2004/005500

PCT/KR2003/001301

The samples were centrifuged at 10,000× g for 5 minutes to obtain supernatants. The supernatants were purified by calmodulin affinity chromatography. Precisely, the supernatants (crude extracts) were applied onto the equilibrated calmodulin affinity chromatography resin, followed by the reaction at 4°C for 24 hours. In order to
5 remove unattached proteins and other substances, the column was washed with CaCl₂ binding buffer. The proteins attached to the calmodulin were isolated from column matrix using elution buffer (50 mM Tris-HCl, pH 8.0, 10 mM β - mercaptoethanol, 2 mM EDTA, 150 mM NaCl). The proteins isolated from crude extracts of *E. coli* transformed with pCAL-n vector were used as a control group.

10 As a result, as seen in FIG. 2, fractions #12-14 of all the fractions separated from *E. coli* transformed with pCKAPA recombinant vector showed the highest protein content. In order to make sure the purity of the proteins, the fractions #12-14 selected from all effluents separated from *E. coli* transformed with pCKAPA recombinant vector and the control group *E. coli*, respectively, were subjected to
15 SDS-PAGE. As a result, as seen in FIG. 3, it was confirmed that the effluents separated from *E. coli* transformed with pCKAPA recombinant vector contain a 55 kDa size fusion protein (51.3 kDa (molecular weight of the protein expressed by *AtKAPAS* gene) + 4 kDa (molecular weight of calmodulin binding peptide)). On the other hand, it was confirmed that the effluents separated from the control group *E.*
20 *coli* do not contain such size protein.

Example 3: Analysis on the enzyme activity of the protein

In order to investigate whether the isolated protein has the function of KAPA syntase that relates to biotin biosynthesis, the present inventors measured the enzyme
25 activity using pimeloyl CoA as a substrate.

WO 2004/005500

PCT/KR2003/001301

The enzyme activity of the protein was measured at 340 nm using a spectrophotometer (Backman DU series 60 spectrophotometer) adjusted at 30°C according to a method of Alexeev *et al.* (Alexeev *et al. J. Mol. Biol.*, 284:401-419, 1998). In the 1 ml of reaction solution for the measurement of the enzyme activity, 20 mM potassium phosphate (pH 7.5), 1 mM NAD⁺, 3 mM MgCl₂, 0.1 unit α-ketoglutarate dehydrogenase and 2-10 μg protein purified in the above Example <2-2> were included. In every reaction solution, the concentration of the protein was 10 μM. L-alanine and pimeroyl CoA were also added into each reaction solution (0-30×10⁸M⁻¹). Results were analyzed using the Soft-Pac Module kinetics software.

Before the analysis, enzyme samples were dialyzed with 20 mM potassium phosphate buffer (pH 7.5) containing 100 μM PLP (pyridoxal 5'-phosphate) at 4°C for 2 hours. The cuvette containing other substances but the purified protein was used as a control group for the measurement of the enzyme activity.

The enzyme activity of the purified protein was suited to Michaelis-Menten kinetics for pimeroyl CoA, a substrate. The changes of reaction speed according to substrate concentration were measured, and then represented by Lineweaver-Burk's plot (see FIG. 4). As a result, K_m value and V_{max} value were 5.4× 10⁻⁷M and 7.93, respectively. Such results supported the fact that the protein expressed by *AtKAPAS* gene is KAPA syntase having substrate specificity to pimeroyl CoA. Also, the K_m values of KAPA syntases of *E. coli* and *Bacillus sphaericus* to pimeroyl CoA were 0.5 mM and 2.5 mM each, while K_m value of the protein of the present invention was 5.4× 10⁻⁷M. Thus, the protein of the present invention was confirmed to have high substrate specificity to pimeroyl CoA, compared with those of *E. coli* and *Bacillus*.

25

WO 2004/005500

PCT/KR2003/001301

Example 4: Preparation transgenic *Arabidopsis thaliana* in which antisense construct of *AtKAPAS* gene was introduced, and its characteristic analysis

<4-1> Preparation of transgenic *Arabidopsis thaliana* in which antisense construct of *AtKAPAS* gene was introduced

- 5 In order to investigate physiological characteristics of the protein purified in the above Example <2-2>, the present inventors prepared transgenic *Arabidopsis thaliana* in which *AtKAPAS* gene was introduced in the antisense orientation, resulting in the inhibition of the expression of *AtKAPAS* transcript.

- The *AtKAPAS* DNA was amplified from cDNA of *Arabidopsis thaliana* by
10 PCR using primers including *Bgl*II site and represented by SEQ. ID. NO. 5 and 6, respectively. The amplified DNA was cut with restriction enzyme *Bgl*II and cloned into pSEN vector prepared to be regulated not by CaMV 35S promoter of pCAMBIA-3301 vector (provided by a laboratory, Department of Life Science, Pohang University of Science and Technology, Korea) but by *sen1* promoter in the
15 antisense orientation in order to avoid the plant death in germination stage, leading to the construction of pSEN-K recombinant vector (see FIG. 5). The *sen1* promoter has specificity to genes expressed in each phase of plant growth.

- The pSEN-K recombinant vector was introduced into *Agrobacterium tumefaciens* using electroporation. The transformed *Agrobacterium tumefaciens* was
20 cultured at 28°C until the OD₆₀₀ reached 1.0. The cells were collected by centrifugation at 5,000rpm for 10 minutes at 25°C. The collected cells were suspended in IM (Infiltration Medium; 1X MS SALTS, 1X B5 vitamin, 5% sucrose, 0.005% Silwet L-77, Lehle Seed, USA) medium until OD₆₀₀ reached 2.0. The 4-week-old *Arabidopsis thaliana* was dipped in *Agrobacterium tumefaciens* suspension
25 in a vacuum chamber and stored under vacuum condition (10⁴ Pa) for 10 minutes. After dipping, *Arabidopsis thaliana* was put in with polyethylene foil for 24 hours.

WO 2004/005500

PCT/KR2003/001301

The transformed *Arabidopsis thaliana* was grown to obtain seeds (T1). For the control groups, wild type *Arabidopsis thaliana* that was not transformed and *Arabidopsis thaliana* transformed with pSEN vector without antisense *AtKAPAS* gene were selected and used.

5

<4-2> Analysis on the characteristics of transgenic *Arabidopsis thaliana* T1

Seeds harvested from the transgenic *Arabidopsis thaliana* obtained in Example <4-1> were selected by dipping it in 0.1% Basta (Kyungnong Co., Korea) solution for 30 minutes and culturing. The transgenic *Arabidopsis thaliana* showed retarded growth or plant death, compared with those of control group (*Arabidopsis thaliana* transformed with pSEN vector).

In order to confirm whether the transgenic plant transformed with antisense construct of *AtKAPAS* gene is a biotin auxotrophic mutant, the reactions induced by adding biotin was investigated. Particularly, 5000 transformed seeds, which were dipped in Basta solution for 30 minutes, were cultured in petri-dishes containing MS medium with or without 5 μ M of biotin. And also, the transformed seeds were cultured in pots filled with sands containing 1 mM of biotin or not. The pots were treated with Basta 5 times, and then investigated for the growth of *Arabidopsis thaliana* in every pot.

As a result, 16 individual plants were grown up in biotin added pots and their phenotype was not much different from that of wild type *Arabidopsis thaliana*. Meanwhile, just 11 individual plants were grown up in pots without biotin. 4 of those showed retarded growth, 2 died and their leaves turned yellow (see FIGs. 6A and 6B). Therefore, the transgenic plants comprising the antisense construct of *AtKAPAS* gene were proved to be biotin auxotrophic mutants.

WO 2004/005500

PCT/KR2003/001301

INDUSTRIAL APPLICABILITY

The present invention provides a polypeptide involved in biotin biosynthesis in plants, a polynucleotide encoding the said polypeptide, a method for inducing
5 plant growth inhibition by suppressing the expression or function of the said polypeptide, resulting in inhibition of biotin biosynthesis, and a method for identifying herbicidal compounds that inhibit the expression or function of the said polypeptide. Since biotin biosynthesis pathway is present neither in human nor in animals, the herbicidal compounds having functions to suppress the expression or
10 function of the polypeptide participating in biotin biosynthesis have an advantage of doing no harm to human and animals. Therefore, such herbicidal compounds can be effectively used as environmental friendly and safe herbicides.

Those skilled in the art will appreciate that the conceptions and specific
15 embodiments disclosed in the foregoing description may be readily utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present invention. Those skilled in the art will also appreciate that such equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended claims.

20

WO 2004/005500

PCT/KR2003/001301

WHAT IS CLAIMED IS:

1. An isolated polypeptide having an amino acid sequence selected from the group consisting of;
5 (a) an amino acid sequence of SEQ. ID. NO. 2, and
(b) an amino acid sequence having at least 70% identity to SEQ. ID. NO. 2.
2. The polypeptide of claim 1, wherein the polypeptide (having the amino acid sequence of SEQ. ID. NO. 2) contains aminotransferase class I and II domain at
10 C-terminal.
3. An isolated polynucleotide encoding the polypeptide of claim 1.
4. The polynucleotide of claim 3, wherein the polynucleotide comprises a nucleotide
15 sequence of SEQ. ID. NO. 1.
5. An expression vector comprising the polynucleotide of claim 3.
6. The expression vector of claim 5, wherein the expression vector comprises the
20 polynucleotide having a nucleotide sequence of SEQ. ID. NO. 1.
7. A cell comprising the expression vector of claim 5.
8. A transgenic plant comprising the expression vector of claim 5.
25

WO 2004/005500

PCT/KR2003/001301

9. A seed comprising the expression vector of claim 5.
10. An antisense polynucleotide complementary to the polynucleotide of claim 3.
- 5 11. The antisense polynucleotide of claim 10, wherein the antisense polynucleotide is complementary to the polynucleotide having a nucleotide sequence of SEQ. ID. NO. 1.
12. An expression vector comprising the antisense polynucleotide of claim 10.
- 10 13. The expression vector of claim 12, wherein the expression vector comprises the antisense polynucleotide of claim 11.
14. A cell comprising the expression vector of claim 12.
- 15 15. A transgenic plant comprising the expression vector of claim 12.
16. A seed comprising the expression vector of claim 12.
- 20 17. A method for inducing plant growth inhibition by suppressing the expression or function of the polypeptide of claim 1, resulting in inhibition of biotin biosynthesis.
- 25 18. The method of claim 17, wherein the suppression of the expression of the polypeptide is achieved by any method selected from the group consisting of introduction of the antisense polynucleotide of claim 10, gene deletion, gene

WO 2004/005500

PCT/KR2003/001301

insertion, T-DNA introduction, homologous recombination and transposon tagging.

19. A method for identifying herbicidal compounds that inhibit the expression or
5 function of the polypeptide of claim 1, comprising the steps of,

(a) combining a polypeptide of claim 1 with the compounds to be tested for the ability to inhibit the expression or the function of the polypeptide under conditions conducive to inhibition;

(b) selecting the identified compounds to inhibit the expression or function
10 of the polypeptide in step (a);

(c) applying the said compounds selected in step (b) to a plant to test for herbicidal activity; and

(d) selecting the identified compounds to have herbicidal activity in step (c).


15

WO 2004/005500

PCT/KR2003/001301

2/7

FIG. 2




WO 2004/005500

PCT/KR2003/001301

3/7

FIG. 3




WO 2004/005500

PCT/KR2003/001301

4/7

FIG. 4




WO 2004/005500

PCT/KR2003/001301

6/7

FIG. 6A




WO 2004/005500

PCT/KR2003/001301

7/7

FIG. 6B



WO 2004/005500

PCT/KR2003/001301

SEQUENCE LISTING

5 <110> GENOMINE INC.
KOREA RESEARCH INSTITUTE OF CHEMICAL TECHNOLOGY

<120> Novel polypeptide having function of 7-keto-8-aminopelargonic acid synthase of plant and method for inducing growth inhibition and lethality by suppressing expression of the polypeptide

10 <130> OP03-1029

<150> KR 2002-38011
<151> 2002-07-02

15 <160> 6

<170> KopatentIn 1.71

20 <210> 1
<211> 1410
<212> DNA
<213> Arabidopsis thaliana

25 <400> 1
atggcggatc attcgtggga taaaactgtg gaagaagcag tgaatgtgct tgaatccagg 60
caaatcttc gatctttgag gccatttgc atgtctaggc aaaacgaaga agaaatagtg 120
30 aaaagcagag ccaatggagg agacgggtac gaggtgttcg acggtttgtg tcaatgggat 180
cggacttcag ttgaggtgtc tgtctgatt cctacatttc agaaatggct tcacgatgaa 240
cccagcaacg gagaagagat ttttagtggg gatgcattag ctgagtgtag aaaagggaga 300
35 ttcaagaagc tgcttttgtt ctctgggaat gattatttgg gtttgagctc acatcctaca 360
atatcaaacg ctgctgcaaa cgcagtcaaa gaatatggta tgggacctaa gggttctgct 420
40 ttaatatgtg gctataccac ttatcatcgt ttgcttgagt ctagtttggc gcaactgaag 480
aaaaaagagg attgtcttgt ttgtcctact gggtttgctg ccaatatggc tgcaatgggt 540
gcaattggaa gtgttgcttc tctttggcc gctagcggga aacctcgaa gaatgaaaaa 600
45 gtggccatct ttctgatgc gctgaatcat gcatcaatta ttgatgtgt cgtcttgct 660
gaacgacaag gaaatgttga agtttttgtt tatcgacact gtgacatatc aaattgcaaa 720

WO 2004/005500

PCT/KR2003/001301

atgaagagga aggtcgtggt gactgatagc ttatttagta tggacggtga cttgcacca 780
 atggaagagc tctctcagct tcggaagagc taaggcctcc ttctagttat tgatgatgct 840
 5 caiggaacat tigtctgtgg agaaaacggt ggtggcgtgg ctgaggaatt taactgtgaa 900
 gctgaigtag attatgtgt gggcactttg agtaaggcag caggggtgtca tggcggtttc 960
 10 atagcttgca gcaaaaaatg gaagcaactg atacagtcca gaggtcgttc attcataatt 1020
 tcaacagcaa tccctgtccc aatggctgca gctgcttaig cagcagtgtg agtggcgagg 1080
 aaggagatat ggagaagaaa ggcaatatgg gagagggtaa aagagttcaa ggaattatct 1140
 15 ggagttgaca tcicaagccc cattatctca ctgtttag ggaatcaaga gaaagccctc 1200
 aaagcgagcc ggtatctatt aaaaicaggc ttccatgtaa tggcaatagc accgcccaca 1260
 20 gtgccacca attctgcag gctaagggtg acactgagtg cagcacatac cacagaagat 1320
 gtgaagaaac tcatcactgc gcttctct tgttggact ttgacaacac agccactcac 1380
 attccttctt ttctatttcc caaattataa 1410
 25
 <210> 2
 <211> 469
 <212> PRT
 30 <213> Arabidopsis thaliana
 <400> 2
 Met Ala Asp His Ser Trp Asp Lys Thr Val Glu Glu Ala Val Asn Val
 1 5 10 15
 35 Leu Glu Ser Arg Gln Ile Leu Arg Ser Leu Arg Pro Ile Cys Met Ser
 20 25 30
 Arg Gln Asn Glu Glu Glu Ile Val Lys Ser Arg Ala Asn Gly Gly Asp
 40 35 40 45
 Gly Tyr Glu Val Phe Asp Gly Leu Cys Gln Trp Asp Arg Thr Ser Val
 50 55 60
 45 Glu Val Ser Val Ser Ile Pro Thr Phe Gln Lys Trp Leu His Asp Glu
 65 70 75 80
 Pro Ser Asn Gly Glu Glu Ile Phe Ser Gly Asp Ala Leu Ala Glu Cys

WO 2004/005500

PCT/KR2003/001301

	85	90	95
	Arg Lys Gly Arg Phe Lys Lys Leu Leu Leu Phe Ser Gly Asn Asp Tyr		
	100	105	110
5	Leu Gly Leu Ser Ser His Pro Thr Ile Ser Asn Ala Ala Ala Asn Ala		
	115	120	125
	Val Lys Glu Tyr Gly Met Gly Pro Lys Gly Ser Ala Leu Ile Cys Gly		
10	130	135	140
	Tyr Thr Thr Tyr His Arg Leu Leu Glu Ser Ser Leu Ala Gln Leu Lys		
	145	150	155
			160
15	Lys Lys Glu Asp Cys Leu Val Cys Pro Thr Gly Phe Ala Ala Asn Met		
	165	170	175
	Ala Ala Met Val Ala Ile Gly Ser Val Ala Ser Leu Leu Ala Ala Ser		
	180	185	190
20	Gly Lys Pro Leu Lys Asn Glu Lys Val Ala Ile Phe Ser Asp Ala Leu		
	195	200	205
	Asn His Ala Ser Ile Ile Asp Gly Val Arg Leu Ala Glu Arg Gln Gly		
25	210	215	220
	Asn Val Glu Val Phe Val Tyr Arg His Cys Asp Ile Ser Asn Cys Lys		
	225	230	235
			240
30	Met Lys Arg Lys Val Val Val Thr Asp Ser Leu Phe Ser Met Asp Gly		
	245	250	255
	Asp Phe Ala Pro Met Glu Glu Leu Ser Gln Leu Arg Lys Lys Tyr Gly		
	260	265	270
35	Phe Leu Leu Val Ile Asp Asp Ala His Gly Thr Phe Val Cys Gly Glu		
	275	280	285
	Asn Gly Gly Gly Val Ala Glu Glu Phe Asn Cys Glu Ala Asp Val Asp		
40	290	295	300
	Leu Cys Val Gly Thr Leu Ser Lys Ala Ala Gly Cys His Gly Gly Phe		
	305	310	315
			320
45	Ile Ala Cys Ser Lys Lys Trp Lys Gln Leu Ile Gln Ser Arg Gly Arg		
	325	330	335
	Ser Phe Ile Phe Ser Thr Ala Ile Pro Val Pro Met Ala Ala Ala Ala		

WO 2004/005500

PCT/KR2003/001301

340 345 350
 Tyr Ala Ala Val Val Val Ala Arg Lys Glu Ile Trp Arg Arg Lys Ala
 355 360 365
 5 Ile Trp Glu Arg Val Lys Glu Phe Lys Glu Leu Ser Gly Val Asp Ile
 370 375 380
 10 Ser Ser Pro Ile Ile Ser Leu Val Val Gly Asn Gln Glu Lys Ala Leu
 385 390 395 400
 Lys Ala Ser Arg Tyr Leu Leu Lys Ser Gly Phe His Val Met Ala Ile
 405 410 415
 15 Arg Pro Pro Thr Val Pro Pro Asn Ser Cys Arg Leu Arg Val Thr Leu
 420 425 430
 Ser Ala Ala His Thr Thr Glu Asp Val Lys Lys Leu Ile Thr Ala Leu
 435 440 445
 20 Ser Ser Cys Leu Asp Phe Asp Asn Thr Ala Thr His Ile Pro Ser Phe
 450 455 460
 25 Leu Phe Pro Lys Leu
 465
 30 <210> 3
 <211> 28
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> Forward primer for AtKAPAS gene
 40 <400> 3
 ggcgatcct tcgcccaaat cacaattc 28
 45 <210> 4
 <211> 32
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Reverse primer for AtKAPAS gene

WO 2004/005500

PCT/KR2003/001301

<400> 4
ggcaagcttt tcactgacaa tatcagaac aa 32

5
<210> 5
<211> 26
<212> DNA
<213> Artificial Sequence

10
<220>
<223> Forward primer for AtKAPAS gene

15 <400> 5
gcagatcttc gcccaaatca caatc 26

20 <210> 6
<211> 31
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Reverse primer for AtKAPAS gene

<400> 6
gcagatcttt cactgacaat atcagaaca a 31

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference OP03-1029	<div style="display: flex; justify-content: space-between;"> <div style="text-align: center;">FOR FURTHER ACTION</div> <div style="font-size: small;">see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.</div> </div>	
International application No. PCT/KR03/01301	International filing date (day/month/year) 02 JULY 2003 (02.07.2003)	(Earliest) Priority Date (day/month/year) 02 JULY 2002 (02.07.2002)
Applicant GENOMINE INC. et al		

This International search report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This international search report consists of a total of 3 sheets.

☐ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the language, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.
- ☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).
- b. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of the sequence listing:
- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☒ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☒ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

2. ☐ Certain claims were found unsearchable (See Box I).

3. ☐ Unity of invention is lacking (See Box II).

4. With regard to the title,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established by this Authority to read as follows:

5. With regard to the abstract,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the drawing to be published with the abstract is Figure No. 6a

- ☒ as suggested by the applicant. ☐ None of the figures
- ☐ because the applicant failed to suggest a figure.
- ☐ because this figure better characterizes the invention.

INTERNATIONAL SEARCH REPORT

National application No.

PCT/KR03/01301

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C12N 9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

-IPC7 C12N 9/10, C12N 15/29, C12N 15/54, C12N 15/82, A01H 5/00, C12N 5/04

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

NCBI PubMed, CA, Esp@cent 7-keto-8-aminopelargonic acid synthase, biotin synthesis, transformed plant, seed

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	GenBank Accession No. T48458 '8-amino-7-oxopelargonic synthase-like protein - Arabidopsis thaliana' 20 April 2000	1-7, 10-14 8, 9, 15, 16
Y	US 5,477,001 A (ELF SANOFI et al.) 19 December 1995 (19.12.1995) See the claims	8, 9, 15, 16
A	RENDINA A.R. et al. The design and synthesis of inhibitors of decthiobiotin synthetase as potential herbicides' In: Pestic Sci, 1999, Vol.55, pp236-247 See the whole document	1-19
A	SCHNEIDER G. & LINDQVIST Y. 'Structural enzymology of biotin biosynthesis' In: FEBS Lett, 2001, Vol.495, pp7-11 See the whole document	1-19
A	PLoux O. & MARQUET A. Mechanistic studies on the 8-amino-7-oxopelargonic synthase, a pyridoxal-5'-phosphate-dependent enzyme involved in biotin biosynthesis' In: Eur J Biochem, 1996, Vol.236, pp301-308 See the whole document	1-19

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

10 OCTOBER 2003 (10.10.2003)

Date of mailing of the international search report

10 OCTOBER 2003 (10.10.2003)

Name and mailing address of the ISA/KR




Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-751,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

KWON, Oh Hye

Telephone No. 82-42-481-5597



INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR03/01301

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5477001 A	19.12.1995	WO 9216632 A1 EP 0536364 A1 JP 6500237 T2	01.10.1992 14.04.1993 13.01.1994

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.